



your community back on track by becoming an EDUCATOR

You can get

SWEDEN 60% GREECE 38%

CANADA 84% SOUTH AFRICA 29%

INDIA 40% **UKRAINE 40%**

BRAZIL 52%

JAMAICA 40%

READ & GET YOUR FACTS STRAIGHT

PRINT & DISTRIBUTE

SHARE ONLINE

TALK ABOUT IT

* (UNATDS, 2010) (UNGASS, CANADA 2012)



- MOSQUITOES DO NOT PASS HIV IF THEY BITE YOU. HIV DOES NOT REPRODUCE OR SURVIVE FOR LONG INSIDE A MOSQUITO.
- HIV IS IN SALIVA, BUT NOT ENOUGH FOR THE VIRUS TO BE TRANSMITTED BY KISSING.
- HIV IS UNABLE TO REPRODUCE OR SURVIVE FOR LONG OUTSIDE OF THE BODY; THERE IS NO RISK FROM TOUCHING SURFACES (TOILET SEATS) OR PEOPLE WITH HIV (HOLDING HANDS).
- HIV CANNOT SURVIVE OR BE TRANSMITTED THROUGH THE AIR (COUGHING) OR WATER (SWIMMING POOLS).



- PREGNANCY & BREASTFEEDING
- JUNPROTECTED VAGINAL, ANAL & ORAL SEX
- RE-USING NEEDLES & SYRINGES TO INJECT DRUGS

34 MILLIGN

Knowing you're positive means you can access excellent treatment

PEOPLE ARE LIVING WITH HIV GLOBALLY*

2-OUT OF 3 PEOPLE ARE UNAWARE OF THEIR HIV-POSITIVE STATUS

Palalalalatatatat

T A T A T A T A T A T A T A T



* (UNAIDS, 2011)

TIP Know your status, GET TESTED!!!



This will enable you to live a full and healthy life



GET TESTED Treating any other STIs that you have reduces the risk that you will contract HIV through sex. You need to get tested before you can get treated.

AT LEAST SO NEW CASES OF STIS PER YEAR* * Based on 4 out f 30 STIS (WHO) 2011

HIV \$ other STI testing is available from health professionals. See 'help \$ advice' at the back.

Health professionals should keep HIV testing confidential.

HTV IS FOUND IN THE SEXUAL FLUIDS OF AN INFECTED MAN OR WOMAN.

THE RISK OF HIV IS HIGHEST WITH PENETRATIVE SEX AND MUCH LOWER WITH . HIV CAN ONLY BE PASSED ON IF HIV-INFECTED BLOOD OR SEXUAL FLUIDS COME INTO CONTACT WITH GENITALS OR A CUT OR SORE IN THE

SALIYA DOES NOT POSE A RISK.



-> USE A CONDOM AND LUBRICANT





If you're HIV-positive discuss different prevention methods with your health professional.

Taking antiretroviral treatment can further reduce the risk of transmission.

A CONDOM

YOU NEED TO USE A SEXUAL CONTACT.

MENTS USE THE SAME CONDOM TWICE.

ONLY PUT ON A CONDOM ONCE THERE IS A PARTIAL OR FULL ERECTION.

TIP If the condom splits: Get a follow-up test for STIs, and consider emergency contraception. If your partner is HIV-positive take PEP. See 'help and advice' at the back of this booklet.







BE CAREFUL NOT TO TEAR THE CONDOM WHEN YOU OPEN IT. CHECK ITS CONDITION AND EXPLRY DATE. PLACE THE ROLLED CONDOM OVER THE TIP OF THE HARD PENIS. PINCH THE TIP AND UNROLL IT. ROLL THE CONDOM
ALL THE WAY DOWN
TO THE BASE OF THE
PENIS, AND SMOOTH
OUT ANY AIR BUBBLES,
WHICH CAN CAUSE A
CONDOM TO BREAK.



Women can also use female condoms. Never use them at the same time as a male condom.

You may find yourself in a situation where your partner doesn't understand how important condoms are ...

Don't you

trust me? I trust you

to make sure we're

both safe.

to do the right thing,

But, we did it before without a condom!

Condoms

ruin the mood.

That was a bad decision. Sex without one puts our health at risk.

I can't

protected.

get in the mood

unless we're both

It doesn't feel as good with a condom on!

Then we need to find a type of condom that can make it feel even better!

Don't you love me? I want to make sure that we're both safe and profected because I love you.

In case of excuses, it's good to get yourself armed with some responses to help

Change their mind!

Reckon you're ready to be an HIV and AIDS Educator? Get your HIV and AIDS knowledge up to scratch with AVERT's quiz:

- 1 What is the difference between HIV and AIDS?
- 2 Can you get AIDS from sharing a cup with someone with HIV?
- 3 Which drugs do people infected with HIV take to prevent the progression to AIDS?
- 4 Complete the sentence: PMTCT stands for Prevention of Mother-to-Child?
- 5 Oral sex is when you talk during sex. True or false?
- 6 Can a woman have a baby if she is HIV-positive?

Complete AVERT's online quizzes to see how you scored!

HELPLINE NUMBERS

AVERT HELP AND INFO SERVICE: (+44) 1403 210 202
AIDSMAP: WWW.AIDSMAP.COM/E-ATLAS
GNP +: (+31) 20 423 4114

HELP WEBSITES

AVERT.ORG: WWW.AVERT.ORG
NAM AIDSMAP: WWW.AIDSMAP.COM
MY HIV CLINIC: WWW.MYHIVCLINIC.COM



HEALTH SERVICES

FIND A CLINIC OFFERING CONTRACEPTION, STI TESTING AND OTHER SEXUAL HEALTH SERVICES.

AIDSMAP: WWW.AIDSMAP.COM/E-ATLAS

POST-EXPOSURE PROPHYLAXI

THIS TREATMENT NEEDS TO BE TAKEN WITHIN 72 HOURS (PREFERABLY 24 HOURS)
TO PREVENT HIV. IF IT'S NOT A WEEKEND OR PUBLIC HOLIDAY, GO TO YOUR
NEAREST SEXUAL HEALTH (GUM) CLINIC. OTHERWISE, VISIT THE
EMERGENCY DEPARTMENT OF YOUR NEAREST HOSPITAL.

USE THIS BOOKLET AND OUR ONLINE RESOURCES TO RAISE AWARENESS AND UNDERSTANDING OF HIV & AIDS.

COMMUNITY PROJECT SUPPORT

AVERT.ORG/AIDS-PROJECTS

WE WORK WITH COMMUNITY-BASED ORGANISATIONS TO RESPOND TO THE HIV & AIDS EPIDEMIC IN SOUTHERN AFRICA

INFORMATION SERVICE

AVERT.ORG/CONTACT

WE CAN PROVIDE YOU WITH INFORMATION, HELP & ADVICE AROUND HIV & AIDS. EMAIL INFO@AVERT. ORG WITH YOUR ENQUIRY.